

Title	Topology of Versal Deformations (超曲面の特異点)
Author(s)	LE, DUNG TRANG
Citation	数理解析研究所講究録 (1975), 237: 55-60
Issue Date	1975-05
URL	http://hdl.handle.net/2433/105523
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Topology of versal deformations.

Lê Dũng Tráng (Ecole Polytechnique, Paris)

We are going to speak about results obtained with the collaboration of B. Iversen and G.M. Greuel.

As these results will be published in [6] and [7] we only write a summary of our lecture.

Let $\varphi: (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^p, 0)$ be a flat holomorphic mapping. ^{then} $(\varphi^{-1}(0), 0)$ is a complete intersection. As G. Fiorina showed ^{in [11]}, we may define $\Phi: (\mathbb{C}^n \times \mathbb{C}^k, 0) \rightarrow (\mathbb{C}^p \times \mathbb{C}^k, 0)$ called the miniversal deformation of $(\varphi^{-1}(0), 0)$. We are going to follow the presentation of Φ as it is done in [10].

As it is explained in [8] and [1], we are interested in computing the local fundamental group of the complement of the discriminant of Δ , discriminant of Φ , in the neighbourhood of the origin 0. Using the results of [3] we obtain that there is a ^{direction of} generic \mathbb{C}^2 -planes of $\mathbb{C}^p \times \mathbb{C}^k$ and a neighbourhood U of 0 in $\mathbb{C}^p \times \mathbb{C}^k$ such that:

- 1) the local fundamental group of the complement of Δ at 0 is given by $U - \Delta$;
- 2) for almost every 2-planes P parallel to the given one, sufficiently near to it and not passing through 0, this local fundamental group is given by $P \cap (U - \Delta)$.

Using results of D. Vohmann in [12] we obtain that for such a 2-plane P , the curve $P \cap \Delta$ has only nodes and cusps as singularities ^{in U} and the number of ^{these} nodes ~~of~~ and cusps depends only on the topology of Δ at 0.

In [7] we compute the number ^{κ} of cusps of $P \cap \Delta$, when $(\varphi^{-1}(0), 0)$ is the germ of a hypersurface with an isolated singularity at 0 and [6] we get this number when $(\varphi^{-1}(0), 0)$ is a general complete intersection. We obtain altogether a formula for the number of nodes which is less satisfying because this formula is hardly computable directly from the equations of $(\varphi^{-1}(0), 0)$.

without using elimination theory.

Now let P_0 be a sufficiently general plane going through the origin in $\mathbb{C}^p \times \mathbb{C}^k$. Then

the set $\Phi^{-1}(P_0 \cap \Delta) \cap C(\Phi)$, where $C(\Phi)$ is the singular locus of Φ is a reduced curve. Let us call $\delta_1 = \dim_{\mathbb{C}} \overline{\mathcal{O}_1} / \mathcal{O}_1$, where \mathcal{O}_1 is the local ring of this curve at 0 , and κ the number of its components. Then we get:

Theorem Let f_1, \dots, f_p be equations of $(\varphi^{-1}(0), 0)$.

Then, replacing eventually f_1, \dots, f_p by linear combination of $u_i f_i$ where u_i is a unit at 0 , we have:

$$\kappa = \mu(\varphi^{-1}(0), 0) - \mu(X', 0) + 2\delta_1 - \kappa$$

where X' is the complete intersection determined by f_1, \dots, f_{p-2} .

In the case of hypersurface we found (cf [7]) :

Corollary $\kappa = \mu + 2\delta_1 - \kappa$

But in this case $\Phi^{-1}(P_0 \cap \Delta) \cap C(\Phi)$ is a complete intersection. In this particular case we can show:

$$\mu_1 = \mu(\Phi^{-1}(P_0 \cap \Delta) \cap C(\Phi), 0) = 2\delta_1 - \kappa + 1$$

Thus: $\kappa = \mu + \mu_1 - 1$

In [7] we show that μ_1 is not a topological invariant and using results of [9] we give a interpretation of the fact that in an analytic family μ and μ_1 are constant .

For instance when $\varphi: (\mathbb{C}^n) \rightarrow (\mathbb{C}, 0)$ is $\varphi(z) = z^n$ then $\kappa = n-2$.

Such results should be complete if one can compute the number of nodes directly from the equations.

Finally a good result should be to relate these computations to the one of the ^{local fundamental group of the} complement of the discriminant Δ at 0. Some recent works of F. Lazzeri • [4], [5] and Gabrielov [2] are going in this direction.

Bibliography.

- [1] E. Brieskorn, Séminaire Bourbaki 1971.
- [2] Gabrielov, in Funk. Anal. i iego Priozhenia, 1972.
- [3] H. Hamm - Lê Dũng Tráng Un théorème de Zariski du type de Lefschetz, Ann. Ec. Norm. Sup. 1973, Paris
- [4] F. Lazzeri, A theorem on the monodromy of isolated singularities, in Singularités à Angers, Asterisque 1974.
- [5] F. Lazzeri - ... , to appear in Inv. Mat.
- [6] Lê Dũng Tráng - G.M. Greuel, to appear.
- [7] Lê Dũng Tráng - B. Iversen, Calcul du nombre de cusps d'une déformation semi-universelle d'une hypersurface complexe, Bull. Soc. Math. Fr., to appear.
- [8] F. Pham, Exposés au Séminaire Leray, Collège de France 1969.

- [9] J. J. Risler Sur les déformations équisingulières
d'idéaux , Bull. Soc. Math. Fr. , t.101, 1973, p.3-16.
- [10] B. Teissier Cycles évanescents, Sections planes et
Conditions de Whitney, in Singularités à Cargèse, Asterisque.
- [11] G. Tiourine : Locally semi-universal flat deformations
... , Math. of the USSR, Izvestia Ak. Nauk, Vol. 3,
N°5, p 967-999, 1969.
- [12] D. Vohmann . Doktor thesis , Bonn, 1973.